



Farm specific transmission patterns of *Fasciola hepatica* in Danish dairy cattle based on different diagnostic methods and monitoring of grazing management

Takeuchi-Storm, Nao; Denwood, Matthew; Petersen, Heidi Huus; Larsen Enemark, Heidi; Thamsborg, Stig Milan

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Takeuchi-Storm, N., Denwood, M., Petersen, H. H., Larsen Enemark, H., & Thamsborg, S. M. (2017). *Farm specific transmission patterns of Fasciola hepatica in Danish dairy cattle based on different diagnostic methods and monitoring of grazing management*. Abstract from 26th International Conference of World Association for the Advancement of Veterinary Parasitology (WAAVP), Kuala Lumpur, Malaysia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



26th International Conference of the World Association for Advancement of Veterinary Parasitology

Theme: ***“Combating Zoonoses: Strength in East-West Partnerships”***

Time: Monday 4 September – Friday 8 September

Venue: Kuala Lumpur Convention Centre, Malaysia

Farm specific transmission patterns of *Fasciola hepatica* in Danish dairy cattle based on different diagnostic methods and monitoring of grazing management

Nao Takeuchi-Storm^{*1}; Matthew Denwood¹; Heidi Huus Petersen²; Heidi Larsen Enemark³; Stig Milan Thamsborg¹

¹Department of Veterinary and Animal Sciences/ University of Copenhagen/ Denmark, ²National Veterinary Institute/ Technical University of Denmark/ Denmark, ³Department of Parasitology/ Norwegian Veterinary Institute/ Norway

A recent survey based on meat inspection data showed that approximately 30% of Danish cattle farms were infected with liver flukes, leading to significant economic losses. Despite the widespread problem, up-to-date knowledge on transmission patterns, diagnostic methods and practical measures for control is still lacking. We therefore initiated a longitudinal, observational study in a few infected dairy farms to elucidate farm specific transmission patterns based on different diagnostic methods and grazing management. Two organic and two conventional dairy farms with high *F. hepatica* antibody levels in bulk tank milk were selected. From each farm a cohort of 40 animals from different age groups (calves, heifers, primiparous and multiparous cows) were sampled 7 times between April 2015 and January 2017. Diagnostic methods included faecal egg count by sedimentation, serum ELISA and coproantigen ELISA. Additionally, monthly bulk tank milk samples were analyzed by ELISA. The analyses are ongoing, but preliminary results indicate that *F. hepatica* is mainly transmitted via summer infection of snails as most animals seroconvert in late autumn without shedding of eggs. However, infection early in the grazing season due to overwintered snails has also been observed. One farm where cows are stabled have had some older cows continuing to shed *F. hepatica* eggs, suggesting long life span of *F. hepatica*, although other routes of infection cannot be ruled out. The final results will provide novel and practical information about different diagnostic tests and transmission patterns related to grazing management on farm-level.

Keywords: *Fasciola hepatica*; cattle; ELISA; bulk tank milk; epidemiology